

group because of the great dynamics in anthropometric characteristics and the corresponding functional parameters (9).

To the best of our knowledge, the functional profile of Romany children has not been a subject of detailed research, which is surprising when considering their vast spatial dissemination. As a result, there is insufficient information regarding the ethnic differences in anthropometric and cardiopulmonary parameters in Romany children.

Subjects and Methods

Subjects and Design

Two hundred and ten healthy children – 153 Bulgarian (75 boys and 78 girls) and 57 Romany (28 boys and 29 girls) of comparable age (mean \pm standard deviation 10.5 ± 0.5 years; range 10-11) took part in the study.

All of the children were pupils of the 4th and 5th grade from a primary school in Plovdiv, South Bulgaria. A total of 250 leaflets of invitation to the trial were distributed among all the pupils from 4th and 5th grade, and we received 217 (85.2%) of them back. Two hundred and ten children completed the study successfully and were taken into account in the statistical processing of the data. The Bulgarian/Romany ratio in the studied group was 3/1 and reflects the real distribution of the two ethnic groups in the respective school (total of 814 pupils – 251 Romany and 563 Bulgarian). The children were in good health, without chronic diseases, taking no medications that might affect exercise capacity. They were generally physically active, but not engaged in regular sports activities. Prior to the test procedures, written informed consent was obtained from a parent or guardian and the associated risks and benefits were explained. The procedures used in this study were approved by the Institutional Ethics Committee of the Plovdiv Medical University.

Anthropometric Measurements

All participants were subjected to complete anthropometric measurements, including skinfolds. The skinfold thicknesses measured by caliper (Harpندن, British Indicators Inc, London, UK) over the triceps and subscapular regions were added together to give the sum of skinfolds and percent body fat (FAT%) was calculated using Slaughter equations (11). The values of body surface area (BSA) were obtained by the method of

Gehan and Georges (12) $BSA = 0.02350 \times Ht^{0.42246} \times Wt^{0.51456}$ (m^2), where Ht is height in cm and Wt is weight in kg. Dimensions of the thorax were assessed through the measurements of chest circumference at full inspiration (CCI) corresponding to total lung capacity (TLC), as well as the measurements of chest circumference at full expiration (CCE), corresponding to residual volume (RV). For stratifying children according to their body mass index (BMI), we used the cut off points published by Cole et al based on BMI centiles for subjects aged 2-18 years (13).

Pulmonary Functional Parameters

Pulmonary function testing was carried out with a diagnostic system MasterScreen Diffusion™ (E. Jaeger, Würzburg, Germany) in a seated position, with a nose clip. We performed spirometry, measurement of diffusion capacity, forced expiration, maximal inspiratory and expiratory pressures.

The equipment was suited for pediatric measurements, ie the chair, mouthpiece, and the whole setting was accommodated to fit children's proportions. Regular calibration was done with a 3 liter calibration syringe.

Forced expiratory maneuvers complied with the general acceptability criteria (14,15). At least three technically acceptable attempts of maximal forced expiratory flow-volume curves were recorded. Spirometric reference values for European children (16), previously validated for Bulgarian population (17), were used.

For the diffusion capacity (transfer factor – $T_{L,CO}$) the mean of two single-breath measurements was used. One hundred seventy one children (127 Bulgarian and 44 Romany) met all the acceptability criteria for the performance of diffusion capacity test. $T_{L,CO}$ and transfer coefficient ($T_{L,CO}/V_A$) values were given unadjusted for hemoglobin (Hb). For the transfer factor, we used reference values developed by us (yet unpublished) on 570 healthy children aged 7-18 years.

The best from at least three measurements for maximal inspiratory (at RV) and expiratory (at TLC) mouth pressure (PI_{max} and PE_{max}) was used for the final calculation.

Cardiopulmonary Exercise Parameters

The treadmill test was performed in a laboratory compliant with the guidelines of the AHA (18). The children were habituated to both

the general environment and the actual procedures. The cardiopulmonary exercise test was carried out on a motor driven, electronically controlled treadmill (TrackMaster™, JAS Fitness Systems, Pensacola, FL, USA) using our modification of the Balke protocol (19), which involved two warm up stages on the level at 2.7 and 4.0 km/h, respectively, and nine one-minute increments with constant velocity of 5.4 km/h, starting from 6% elevation and increasing with 2% every minute until exhaustion or elevation of 22%. Recovery period had standard three minutes duration (2.7 km/h and zero elevation).

Throughout the test, gas exchange variables were determined with an on-line computerized system CardiO₂™ (Medical Graphics, St Paul, MN, USA) using standard open circuit techniques. Data were averaged every 30 s and used to calculate oxygen uptake (VO₂; mL/min, standard temperature pressure dry, STPD), and carbon dioxide output (VCO₂; mL/min, STPD and respiratory exchange ratio, RER). The system was calibrated before each test with gases of known concentrations. Heart rate was monitored electrocardiographically (Hellige, Freiburg, Germany) and the oxygen saturation was traced with pulseoxymeter Pulseox DP-8 (Minolta, Osaka, Japan).

Anaerobic threshold (AT) was determined as the level of VO₂ at which at least one of the following was present: 1) increase in ventilatory equivalents for oxygen (V_E/VO₂) without simultaneous increase in ventilatory equivalents for carbon dioxide (V_E/VCO₂); and 2) disappearance of the linear relation between VCO₂ and VO₂ (V-slope method).

Other Measurements

At the end of each exercise increment and throughout the recovery period the children were asked to rate the perceived exertion using the Borg Category-Ratio Scale (20) depicting fatigue

(dyspnea) from “not at all” to “maximal” by means of ten grades.

For clarifying children's health and social status, the parents were asked to fill out a questionnaire on child's birth weight and gestational age (estimated by the calendar method), disease history since birth, the presence of symptoms and complaints for the last 12 months, rate of physical activity, presence of pets at home, domestic hygienic conditions, type of domestic heating, smoking at home, family social status, and number of children in the family.

Statistical Methods

All values were expressed as mean ± standard deviation. The distribution of quantitative variables was tested by the Kolmogorov-Smirnov test. The following statistical analyses were performed: descriptive statistics, Mann-Whitney or Student *t* test where appropriate, χ^2 test, Fisher exact test, correlation analysis, and stepwise regression analysis. All analyses were performed using the Statistical Package for Social Sciences Package for Windows, version 10.05 (SPSS Inc., Chicago, IL, USA). A level of *P* < 0.05 was considered to be statistically significant.

Results

The anthropometric measurements in Bulgarian and Romany children showed that Bulgarian children were significantly taller and heavier in comparison with Romany children (Table 1).

They also differed significantly with regard to the amount of fat. The overweight children constituted 22.2% of the Bulgarian group (7.8% with obesity) whereas there were 14.0% (1.8% with obesity) in the group of Romany children (*P* = 0.516). Significant differences existed only in the measured chest circumference at full inspiration.

The main pulmonary function parameters are presented in Table 2. The mean absolute

Table 1. Anthropometric parameters (mean ± standard deviation) in Bulgarian and Romany children, Bulgaria 2004

Parameters	Bulgarian children	Romany children	Z value*	P
Age (years)	10.5 ± 0.5	10.5 ± 0.5	-0.13	0.898
Height (cm)	145.1 ± 6.7	141.8 ± 6.6	-3.02	0.003
Weight (kg)	38.7 ± 9.2	35.2 ± 7.8	-2.80	0.005
Body mass index (kg/m ²)	18.2 ± 3.3	17.4 ± 2.8	-1.55	0.122
Body surface area (m ²)	1.26 ± 0.17	1.18 ± 0.14	-3.04	0.002
Fat (%)	20.5 ± 8.1	17.8 ± 7.6	-2.32	0.020
Chest circumference at full inspiration (cm)	73.4 ± 6.1	71.9 ± 6.2	-2.09	0.036
Chest circumference at full expiration (cm)	68.1 ± 6.8	67.1 ± 6.1	-0.929	0.353

*Mann-Whitney *U* test.

Table 2. Pulmonary function parameters (mean±standard deviation) in Bulgarian (n=127) and Romany (n=44) children, Bulgaria 2004

Parameter*	Bulgarian children	Romany children	Z† value	P
VC (mL) (% predicted)	2,598±378 (99.2±10.3)	2,396±372 (97.3±11.0)		
FEV ₁ (mL) (% predicted)	2,290±334 (106.3±12.1)	2,123±338 (105.0±11.3)	-3.30 (-0.65)	0.001 (0.514)
FEV ₁ /VC (%)	88.3±5.8	88.8±5.8	-0.62	0.534
PI _{max} (cm H ₂ O)	86±20	81±18	-1.82	0.069
PE _{max} (cm H ₂ O)	114±24	110±21	-0.93	0.354
T _{LCO} † (mmol×min ⁻¹ ×kPa ⁻¹) (% predicted)	5.75±1.0 (96.7±12.7)	5.16±0.96 (92.4±11.0)	-3.78 (-2.65)	<0.001 (0.008)
T _{LCO} /V _A (mmol×min ⁻¹ ×kPa ⁻¹ ×L ⁻¹)	1.69±0.20	1.68±0.22	-0.07	0.946

*Abbreviations: VC – vital capacity; FEV₁ – forced expiratory volume in 1 s; PI_{max} – maximal inspiratory pressure; T_{LCO} – carbon monoxide diffusion capacity (transfer factor); T_{LCO}/V_A – transfer coefficient.

†Mann-Whitney U test.

Table 3. Cardiopulmonary parameters (mean±standard deviation) in Bulgarian (n=127) and Romany children (n=44), Bulgaria 2004

Parameter*	Bulgarian children	Romany children	Z value†	P
V _T (mL)	940±213	889±201	-1.71	0.087
Breathing frequency (min ⁻¹)	45.5±9.2	46.0±8.3	-0.56	0.575
V _E (L/min)	42.0±10.2	40.3±9.8	-1.19	0.235
VO ₂ peak (mL/min)	1341±293	1260±280	-1.92	0.054
VO ₂ /kg (mL×kg ⁻¹ ×min ⁻¹)	34.9±3.7	36.1±5.2	-1.00	0.319
VO ₂ /BSA (mL×min ⁻¹ ×m ⁻²)	1059±124	1059±150	-0.38	0.705
VO ₂ /HR (mL×min ⁻¹ ×beat ⁻¹)	7.5±1.7	7.6±1.7	-0.20	0.843
Heart rate (beat/min)	179±9	167±12	-6.06	<0.001
Respiratory exchange ratio	1.08±0.08	1.08±0.04	-0.42	0.674
V _E /VCO ₂ AT	29.0±2.1	29.7±1.8	-1.39	0.211
Borg score	4.9±1.5	3.2±1.3	-6.43	<0.001

*Abbreviations: V_T – tidal volume; V_E – minute ventilation; VO₂ peak – peak oxygen uptake; BSA – body surface area; VO₂/HR – oxygen pulse; V_E/VO₂ – ventilatory equivalent for oxygen; V_E/VCO₂ – ventilatory equivalent for carbon dioxide; AT – anaerobic threshold.

†Mann-Whitney U test.

values for vital capacity (VC), forced expiratory volume in 1 s (FEV₁), and T_{LCO} were respectively 7.8%, 7.3%, and 10.2% lower in Romany children compared with Bulgarian. No differences were detected with regard to PI_{max} and PE_{max}. There were no differences between Bulgarian and Romany children concerning the volume-standardized parameters – forced expiratory volume in 1 s/vital capacity (FEV₁/VC %) and T_{LCO}/V_A. With the exception of T_{LCO} there were not significant differences in percent predicted values.

The results from the cardiopulmonary exercise test in the studied groups are presented in Table 3. There were no significant differences between the two ethnic groups regarding ventilatory variables (tidal volume (V_T), breathing frequency (BF), and minute ventilation (V_E)), V_E/VO₂, equivalents for carbon dioxide at anaerobic threshold (V_E/VCO₂AT), VO₂ at peak exercise (VO₂peak), as well as in its “normalized” derivatives (oxygen uptake per kg (VO₂/kg), oxygen uptake/body surface area (VO₂/BSA) and oxygen pulse (VO₂/HR)). It is worthy to note the significantly higher heart rate in Bulgarian children. The level of physical exertion is almost the same in the studied groups (comparable RER) but the perception of exertion was significantly higher in Bulgarian children as seen from the Borg score.

Only two Romany children (3.5%) did not complete the test whereas 12 (7.8%) Bulgarians discontinued the standardized protocol prematurely (P=0.423). Eleven of these children appeared to be overweight or obese according to the cut off values of Cole et al (13). Children who did not complete the test (n=14) had significantly higher Borg score compared to the rest of the subjects (7.5 ± 0.7 vs 4.2 ± 1.5; P<0.001).

Significant gender differences were established simultaneously in two ethnic groups with regard to the following parameters: V_E/VO₂ and V_E/VCO₂AT (higher in girls), PI_{max} and PE_{max} (higher in boys), and FAT% (higher in girls). Romany boys had significantly higher values for VO₂ peak and its derivatives (VO₂/kg, VO₂/HR), whereas for Bulgarians had significantly higher lung function parameters (VC, FEV₁ and T_{LCO}).

A box-plot of birth weight of 83 Bulgarian (45 boys/38 girls) and 41 Romany (20 boys/21 girls) children from the whole group is presented in Figure 1. We took into consideration only those infants delivered in term whose birth weight and gestational age were known. Bulgarian children had significantly higher birth weight in comparison with Romanies (3,358 ± 513 vs 3,095 ± 435 g; P=0.006). The children's birth weight correlated

Table 4. Determinants of the main studied parameters and equations characteristics*

Variable	Predictors	β -coefficient	SE	t	P	Model characteristics
						F, significance
Vital capacity	constant	-3.159	0.469	-6.74	<0.001	88.0, <0.001
	height	0.041	0.003	12.79	<0.001	
	gender	-0.160	0.045	-3.54	0.001	
FEV ₁	constant	-2.613	0.362	7.21	<0.001	99.2, <0.001
	height	0.034	0.002	-13.89	<0.001	
	gender	-0.076	0.034	-2.25	0.026	
T _{LCO}	constant	-7.103	2.067	-3.44	0.001	35.7, <0.001
	height	0.094	0.014	6.68	<0.001	
	parents' smoking	-0.737	0.172	-4.28	<0.001	
VO ₂ peak	constant	260.422	44.049	5.91	<0.001	609.8, <0.001
	weight	28.033	1.135	24.70	<0.001	

*Abbreviations: SE - standard error of β -coefficient; FEV₁ - forced expiratory volume in 1 s; T_{LCO} - carbon monoxide diffusion capacity (transfer factor).




Figure 1. Box-plot of birthweight of Bulgarian and Romany newborns. The box represents the interquartile range which contains the 50% of values. The whiskers are bars that extend from the box to the highest and lowest values, excluding outliers. The line across the box indicates the median.

significantly and positively with VC ($r=0.299$; $P=0.001$), FEV₁ ($r=0.229$; $P=0.010$), T_{LCO} ($r=0.375$; $P<0.001$), VO₂ peak ($r=0.280$; $P=0.012$), as well as with the main anthropometric parameters – height ($r=0.379$; $P<0.001$), weight ($r=0.298$; $P=0.001$), BSA ($r=0.333$; $P<0.001$), and CCI ($r=0.263$; $P=0.003$). However, following application of stepwise regression analysis it became apparent that despite these significant correlations, birth weight was not a determinant of any of the main cardiopulmonary parameters (Table 4). Table 4 represents the determinants of the main studied parameters and equation characteristics.

It is clearly seen from Table 4 that height and gender are the main determinants of children's lung function indices. Body weight, but not

height, was identified as a main determinant of peak exercise oxygen uptake (VO₂ peak). According to this analysis, parents' smoking has a significant effect on the transfer factor. It is worthy to note the absence of ethnic belonging as a determinant in these equations.

We found a higher prevalence of indoor smoking by parents of Romanies in comparison with Bulgarians (59.6% vs 17.6%, respectively; $P<0.001$). Another important environmental concern is the use of fossil fuels for heating without adequate ventilation in the majority of Romany homes (64.9% vs 23.5% in Bulgarian; $P<0.001$).

Discussion

This study showed that there were significant differences between Bulgarian and Romany children with regard to anthropometric parameters, including birth weight, lung function indices, and perception of exertion. The children's birth weight correlated significantly with main anthropometric and cardiopulmonary parameters, but it was not a determinant of the latter.

The limitations of our study were that we did not measure some specific anthropometric variables, such as sitting height and all thoracic dimensions, and we cannot comment on different body proportions, such as trunk/leg length ratio. Yet, according to Cotes (9), this ratio is similar in Caucasians and Indian subjects.

Another limitation of this trial was that the investigation was cross-sectional and covered only a narrow age interval. Despite the imperfections of the study, the results convincingly indicated that stated differences between Bulgarian and Romany children really existed and were not a mere result of differences in body proportions.

According to Crapo et al (7), the best way to document ethnic differences is to confront the functional parameters obtained in the frame of one trial performed by means of standardized methods and design, applied by one and the same team of researchers to the different ethnic groups. Our study was fully compliant with these recommendations and the children were of the same age, so we assume that stated differences really exist. The core of the problem is that, besides genetic factors, some other factors, such as environmental and economic factors could significantly contribute to these differences, and it is difficult to weigh precisely the impact of each of them.

Data in the literature prove that racial variations determining genetic differences in anthropometric characteristics lead to significant differences in lung functional parameters (9,21). Significantly lower values of anthropometric and lung function parameters, found in the Romany children, coincide with the findings of lower values of the same parameters in the North Indian population which is the origin of the Romanies, compared with the white population (2). Yang et al (22) found that the values of lung function parameters for Caucasians are higher than those for Chinese and Indian populations of all ages and heights. Gourgoulianis et al (23) found the lower values of lung function parameters in Gypsies aged 14-70 years in Greece. Having in mind these facts, we have to assume that lower anthropometric and absolute spirometric values of Romany children did not result from some kind of physical retardation but represented at least partially genetically determined lower anthropometric characteristics, among which are stature and body mass.

The results of this study and the data in the literature (22,23) suggest that specific reference values should be used for lung function parameters of Romany population. The general reference values for spirometric measurements in Caucasians can be used for Romany children in case they are built solely on anthropometric variables and not age. The lack of differences in percent predicted spirometric values suggests that standing height is a proxy for lung dimensions in this very narrow age interval (10-11 years). Still, there is general agreement that parameters of lung growth during childhood and adolescence are not determined only by height. Theoretically, not only height and age, but also stages of pubertal devel-

opment should be included in the predictions to improve the quality of pediatric pulmonary reference equations (24).

Despite the genetic factors determining stature and body mass, another reason for lower anthropometric parameters in Romany population can be undernourishment determined by their poor economic status, low educational level, and unsettled lifestyle (23). We found that 4 of the Romanies (7%) had BMI values below the 90th centile according to Cole et al (13), being underweight and presumably inadequately nourished (25). In addition, the percentage of fat in Romany children was significantly lower in comparison with Bulgarians.

Harik-Khan et al (26) point out that factors probably accounting for the racial or ethnic difference may be prenatal or childhood factors such as low birth weight, maternal smoking during pregnancy, allergy, indoor air pollution, and exposure to environmental toxins. The majority of these factors are also found in Romanies.

Based on the questionnaire data obtained by the participants, we can indicate the poorer housing and the use of fossil fuels for heating without adequate ventilation in the majority of Romanies' homes. There was also a higher prevalence of indoor smoking by their parents. In another study we found that more Romany women continued to smoke during pregnancy, compared with Bulgarian women (27). The Romany children in our sample were born by very young mothers (mean age 19.5 ± 3.1 years) which is a consistent sign of social immaturity. Higher degree of indoor air pollution, including exposure to tobacco smoke, in Romanies' homes may at least partially explain significantly lower values (absolute and percent predicted) of their diffusion capacity.

Our results also showed that Romany mothers gave birth to babies with lower birth weight in comparison with Bulgarian mothers. There is a consensus about Romany babies being consistently smaller, although the data on this problem are scarce (28,29). On the other hand, there is a lot of conflicting evidence about the impact of birth weight on health status and lung function parameters (30). We found that birth weight correlated significantly with anthropometric characteristics and some cardiopulmonary parameters, which corresponds to the findings of Rona et al (31).

It is difficult to interpret the significant difference in perceived exertion between the two groups because of a lack of similar data in the literature. A possible explanation is that Romany children who live in harder socioeconomic conditions had higher tolerance towards physical exertion and stress. Ethnic differences are described with regard to the perception of angina pectoris during exercise (32) and perception of dyspnea evoked by bronchial challenge tests (33) but in trials including only adult population.

In line with Cotes (9), who considered exercise parameters as independent of ethnic group, we found that VO_2 peak and its standardized indices VO_2/kg and VO_2/BSA did not differ between Romany and Bulgarian children. Trowbridge et al (34) found that there were no ethnic differences only with regard to resting and submaximal VO_2 , and they are apparent for VO_2 peak. The finding that 96.5% of the Romany children completed the standardized test (vs 92.2% in Bulgarian), their significantly lower score of perception to exercise, and especially significantly lower heart rate, suggest that their functional capacity might be even better than that of their Bulgarian counterparts.

In conclusion, there were significant differences between Bulgarian and Romany children with regard to anthropometric parameters including birth weight, lung function indices, and perception of exertion. These differences appear to be a combined result of biological and environmental factors.

References

- McKee M. The health of gypsies. *BMJ*. 1997;315:1172-3.
- Ginter E, Krajcovicova-Kudlackova M, Kacala O, Kovacic V, Valachovicova M. Health status of Romanians (Gypsies) in the Slovak Republic and in the neighbouring countries. *Bratisl Lek Listy*. 2001;102:479-84.
- Hajioff S, McKee M. The health of the Roma people: a review of the published literature. *J Epidemiol Community Health*. 2000;54:864-9.
- Abel-Smith B, Figueras J, Holland W, McKee M, Mossialos E. Choices in health policy: an agenda for the European Union. Luxembourg: Office for Official Publications of the European Communities; 1995.
- Schwartz J, Katz SA, Fegley RW, Tockman MS. Sex and race differences in the development of lung function. *Am Rev Respir Dis*. 1988;138:1415-21.
- Roizin H, Szeinberg A, Tabachnik E, Molho M, Benzaray S, Augarten A, et al. Ethnic differences in lung function in Israeli children. *Thorax*. 1993;48:906-10.
- Crapo RO, Jensen RL, Oyunchimeg M, Tsh T, DuWayne Schmidt C. Differences in spirometry reference values: a statistical comparison of a Mongolian and a Caucasian study. *Eur Respir J*. 1999;13:606-9.
- Korotzer B, Ong S, Hansen JE. Ethnic differences in pulmonary function in healthy nonsmoking Asian-Americans and European-Americans. *Am J Respir Crit Care Med*. 2000;161(4 Pt 1):1101-8.
- Cotes JE. Lung function: assessment and application in medicine. 5th edition. London: Blackwell Scientific Publications; 1993.
- American Thoracic Society. Lung function testing: selection of reference values and interpretative strategies. *Am Rev Respir Dis*. 1991;144:1202-18.
- Slaughter MH, Lohman TG, Boileau RA, Horswill CA, Stillman RJ, Van Loan MD, et al. Skinfold equations for estimation of body fatness in children and youth. *Hum Biol*. 1988;60:709-23.
- Gehan ES, Georges S. Estimation of human body surface area from height and weight. *Cancer Chemotherapy Report*. 1970;54:225-35.
- Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ*. 2000;320:1240-3.
- American Thoracic Society. Standardization of spirometry, 1994 update. *Am J Respir Crit Care Med*. 1995;152:1107-36.
- European Respiratory Society. Standardized lung function testing. Lung volumes and forced ventilatory flows. *Eur Resp J*. 1993;16 Suppl 6:5-40.
- Quanjer PH, Borsboom GJ, Brunekreff B, Zach M, Forche G, Cotes JE, et al. Spirometric reference values for white European children and adolescents: Polgar revisited. *Pediatr Pulmonol*. 1995;19:135-42.
- Kostianev S, Iluchev D. Is the problem concerning the children's reference spirometric values in Europe solved? [in Bulgarian] *Pediatrics*. 1996;36:37-8.
- Pina IL, Balady GJ, Hanson P, Labovitz AJ, Madonna DW, Myers J. Guidelines for clinical exercise testing laboratories. A statement for healthcare professionals from the Committee on Exercise and Cardiac Rehabilitation, American Heart Association. *Circulation*. 1995;91:912-21.
- Marinov B, Kostianev S, Turnovska T. Submaximal treadmill test for screening of physical capacity in pediatric age group [in Bulgarian]. *Pediatrics*. 2000;40:37-8.
- Borg GA. Psychophysical bases of perceived exertion. *Med Sci Sports Exerc*. 1982;14:377-81.
- Donnelly PM, Yang TS, Peat JK, Woolcock AJ. What factors explain racial differences in lung volumes? *Eur Respir J*. 1991;4:829-38.
- Yang TS, Peat J, Keena V, Donnelly P, Unger W, Woolcock A. A review of the racial differences in the lung function of normal Caucasian, Chinese and Indian subjects. *Eur Respir J*. 1991;4:872-80.
- Gourgoulialis KI, Tsoutsou P, Fotiadou N, Samaras K, Dakis D, Molyvdas PA. Lung function in Gypsies in Greece. *Arch Environ Health*. 2000;55:453-4.
- Merkus PJ, Tiddens HA, de Jongste JC. Annual lung function changes in young patients with chronic lung disease. *Eur Respir J*. 2002;19:886-91.
- Ong TJ, Mehta A, Ogston S, Mukhopadhyay S. Prediction of lung function in the inadequately nourished. *Arch Dis Child*. 1998;79:18-21.

- 26 Harik-Khan RI, Fleg JL, Muller DC, Wise RA. The effect of anthropometric and socioeconomic factors on the racial difference in lung function. *Am J Respir Crit Care Med.* 2001;164:1647-54.
- 27 Kostianev S, Karakashian M. Tobacco smoking (up to 10 cigarettes daily) and lung functions in pregnant women. *Folia Med (Plovdiv).* 1994;36:21-5.
- 28 Adrai J, Ruf H, Blanc B, Papahadjji M, Dupont S, Conte M, et al. Pregnancy and childbirth in gypsies [in French]. *Rev Fr Gynecol Obstet.* 1985;80:679-87.
- 29 Joubert K. Size at birth and some sociodemographic factors in gypsies in Hungary. *J Biosoc Sci.* 1991;23:39-47.
- 30 Edwards CA, Osman LM, Godden DJ, Campbell DM, Douglas JG. Relationship between birth weight and adult lung function: controlling for maternal factors. *Thorax.* 2003;58:1061-5.
- 31 Rona RJ, Gulliford MC, Chinn S. Effects of prematurity and intrauterine growth on respiratory health and lung function in childhood. *BMJ.* 1993;306:817-20.
- 32 Sheffield D, Kirby DS, Biles PL, Sheps DS. Comparison of perception of angina pectoris during exercise testing in African-Americans versus Caucasians. *Am J Cardiol.* 1999;83:106-8, A8.
- 33 Hardie GE, Janson S, Gold WM, Carrieri-Kohlman V, Boushey HA. Ethnic differences: word descriptors used by African-American and white asthma patients during induced bronchoconstriction. *Chest.* 2000;117:935-43.
- 34 Trowbridge CA, Gower BA, Nagy TR, Hunter GR, Treuth MS, Goran MI. Maximal aerobic capacity in African-American and Caucasian prepubertal children. *Am J Physiol.* 1997;273(4 Pt 1):E809-14.

Received: October 28, 2004

Accepted: March 1, 2005

Correspondence to:

Stoilka Mandazhieva
Department of Pathophysiology
Medical University of Plovdiv
15A Vassil Aprilov Blvd.
4002 Plovdiv, Bulgaria
stoilka@mail.orbitel.bg